

Pharmacological potential of an important medicinal plant *Phyllanthus amarus* Schum. & Thonn.

S. S. Sravanthi Pammi, T. V. S. Padmavathi and Archana Giri*

Centre for Biotechnology, Institute of Science and Technology, Jawaharlal Nehru Technological University Hyderabad, Kukatpally, Hyderabad 500 085, India.

*Corresponding Author's Email: archanagirin@yahoo.co.in

ARTICLE INFO

Article history:

Received : 23 May 2016
Accepted : 30 June 2016
Available online : 20 July 2016

Keywords:

Phyllanthus amarus,
traditional uses,
chemical constituents,
pharmacological properties.

ABSTRACT

Herbs have always formed an integral part of human health and are used in the treatment of several human diseases. One of such species which have wide patronage of use is *Phyllanthus amarus*. This herb is in traditional medicine for more than 3,000 years. The plant has also served as lead for several experimental investigations that explored its phytochemical constituents and pharmacological uses. Present paper compiles traditional uses, phytoconstituents and pharmacological properties of *Phyllanthus amarus*.

© 2016 International Journal of Advanced Research in Science and Technology (IJARST).

All rights reserved.

PAPER-QR CODE



Volume 5, Issue 2

Citation: S. S. Sravanthi Pammi et. al., Pharmacological potential of an important medicinal plant *Phyllanthus amarus* Schum. & Thonn, Int. J. Adv. Res. Sci. Technol. Volume 5, Issue 2, 2016, pp.577-582.

Introduction:

Phyllanthus amarus is a plant of the family Euphorbiaceae and has about approximately 800 species which are found in tropical and subtropical countries of the world [1, 2]. The name '*Phyllanthus*' means "leaf and flower" and named so because of its appearance where flower, fruit and leaf appears fused [3]. *Phyllanthus amarus* is a branching annual glabrous herb which is 30-60 cm high and have slender, leaf-bearing branchlet, distichous leaves which are sessile elliptic-oblong, obtuse, rounded base. Flowers are yellowish, whitish or greenish, axillary, male flowers in groups of 1-3 whereas females are solitary. Fruits are depressed-globose like smooth capsules present underneath the branches and seeds are trigonous, pale brown with longitudinal parallel ribs on the back [4]. The plant has been found in Philippine, Cuba, Nigeria and among others. In India, *Phyllanthus amarus* is widely distributed as a weed in cultivated and waste lands [5].

Traditional Uses:

Phyllanthus amarus herb has found its traditional usefulness in several health problems such as diarrhoea, dysentery, dropsy, jaundice, intermittent fevers,

urinogenital disorders, scabies and wounds. Further, these are used in the treatment of kidney problems, urinary bladder disturbances, pain, gonorrhea, diabetes and chronic dysentery. Topically, it is used for several skin problems ranging from skin ulcers, sores, swelling and itchiness, wounds, bruises, scabies, ulcers and sores, edematous swellings, tubercular ulcers, ringworm, scabby and crusty lesions. Its effect in excretory system is due to its antiurolithic property and is used in the treatment of kidney/gallstones, other kidney related problems, appendix inflammation and prostate problems [6-8]. Because of its efficacy in the field of gastro-intestinal disorders it is used in the treatment of disorders like dyspepsia, colic, diarrhea, constipation and dysentery. The herb has found use in several female problems such as in leucorrhoea, menorrhagia and mammary abscess and can act as galactagogue. The young shoots of plant are administered in the form of an infusion for the treatment of chronic dysentery. Fresh leaf paste has wound healing capacity and used to cure white spots on skin & jaundice. The stem juice is also used as wound healers. The whole plant extract is used in urinary problems & swelling of liver. The root extract is used to cure stomach pain. The flower paste of plant is applied externally as antidote against snake bite [9-15].

Phytochemical Studies:

Phyllanthus amarus have numerous phytochemicals such as alkaloids, flavonoids, tannins, lignins, polyphenolic compounds and tetracyclic triterpenoids. Several phytoconstituents isolated from this plant are enlisted in Table 1.

Table: 1.Phytochemicals in *Phyllanthusamarus*.

Alkaloids	Isobubbialine and Epibubbialine [16]
Tannins	Geraniin, corilagin, 1,6-digalloylglucopyranoside rutin , quercetin-3-O-glucopyranoside , Amarulone, Phyllanthusiin D &Amariin [17, 18] .
Lignans	Niranthin, Nirtetralin, Phyltetralin, Hypophyllanthin, Phyllanthin, demethylenedioxy-niranthin, 5-demethoxy-niranthin, Isolintetralin [19, 20, 21, 22].
Ellagitannins	Amariin, 1-galloyl-2, 3-dehydrohexahydroxydiphenyl (DHHDP) - glucose, Repandusinic acid, Geraniin, Corilagin, Phyllanthusiin D, and flavonoids namely rutin, and quercetin 3-O-glucoside, 1-Ogalloyl-2,4-dehydrohexahydroxydiphenoyl-glucopyranose elaeocarpusin, repandusinic acid A and geraniinic acid [23, 24] .
Volatile oil	Linalool and Phytol [25].
Triterpene	(2Z, 6Z, 10Z, 14E 18E, 22E-farnesil farnesol) [22].

Pharmacological Activity:

Anticancer activity:

The aqueous extract of *Phyllanthus amarus* demonstrates potent anticancer activity against 20-methylcholanthrene (20-MC) induced sarcoma development. The aqueous extract inhibits DNA topoisomerase II of mutant cell cultures and inhibited cell cycle regulatory enzyme cdc 25 tyrosine phosphatase of *Saccharomyces cerevisiae*. The anticarcinogenic and anti-tumour activity of *Phyllanthus amarus* proposed to be inhibition of metabolic activation of carcinogen as well as the inhibition of cell cycle regulators responsible for cancerous growth and DNA repair [26].

Antiamnesic Activity:

Antiamnesic activity of aqueous extract of leaves and stems of *Phyllanthus amarus* were evaluated for nootropic effects and brain cholinesterase activity in male Swiss albino mice. Scopolamine and diazepam were used as standard drugs to produce amnesia and elevated plus maze and passive avoidance paradigm as models for evaluation of cognitive functions. The result reveals a dose dependent attenuation of diazepam and scopolamine induced amnesic deficits and reduction in brain cholinesterase activity. Since the reduction in cholinesterase is linked with increase acetylcholine concentration in brain which further is responsible for

improving memory, provide a rationale to use this therapeutic potential in the management of patients with cognitive disorders [27].

Antioxidative Activity

The DPPH assay is used to determine antioxidant potential, which is based on the reduction of stable radical DPPH to yellow coloured diphenylpicryl hydrazine. Thus, the ability of the test samples to quench this radical is a measure of its antioxidative ability. *Phyllanthus amarus* have powerful antioxidant property which is evident from the present study in which phyllanthin and *Phyllanthus amarus* extract were evaluated. In the experiment, it was observed that the DPPH free radical scavenging activity was concentration dependent and reaches maximum at a concentration of 20 mol/ml for phyllanthin and 300 g/ml for *Phyllanthus amarus* extract. Further, since phyllanthin possess very high antioxidative property as evident by its low IC₅₀ value of 7.4 mol/ml as compared to *Phyllanthus amarus* extracts suggest its contribution in antioxidative effects [7]. In another study, it has been found that boiled water extract of the fresh and dried *Phyllanthus amarus* plant had comparatively greater antioxidant activity than microwave assisted extraction method employed for the extraction [28].

Antinociceptive Activity:

The hydroalcoholic extract of four *Phyllanthus* species namely *Phyllanthus amarus*, *Phyllanthus orbiculatus*, *Phyllanthus fraternus* and *Phyllanthus stipulatus* were given intraperitoneally and evaluated in acetic acid-induced writhing and formalin and capsaicin-induced licking effects. In acetic acid-induced writhing test it was found that all produced significant inhibition of acetic acid-induced abdominal constrictions, with mean ID₅₀ values of 0.3, 1.8, 7.4 and 26.5 mg/kg for *Phyllanthus amarus*, *Phyllanthus orbiculatus*, *Phyllanthus fraternus* and *Phyllanthus stipulatus*, respectively. Similarly, in the formalin test, it was observed that the hydroalcoholic extract of four species produced graded inhibition against both phases of formalin-induced licking, inhibition in licking being more active in the late phase. Apart from the above models, hydroalcoholic extract of the species also elicited significant reduction in the capsaicin-induced neurogenic pain. It was also observed that hydroalcoholic extract of the *Phyllanthus* species was less potent and efficacious when given orally compared to intraperitoneal route [4].

Antimicrobial activity:

Antimicrobial activity of ethanol and water extracts of *Phyllanthus amarus* were evaluated against the test organisms *Salmonella typhi*. Ethanolic, cold water extract and hot water extract of *Phyllanthus amarus* were employed for antimicrobial evaluation by agar cup diffusion method which were compared against standard antibiotics that were evaluated by disk

diffusion method. The result demonstrates ethanolic extract to be most potent against the test bacteria with diameter of 8.0 mm as growth inhibition zone. This study establishes one of the traditional uses of *Phyllanthus amarus* against typhoid fever [29]. In another study, hexane, petroleum ether, chloroform, acetone and methanol extract of *Phyllanthus* leaves were tested for antibacterial activity against *Pseudomonas aeruginosa*, *Klebsiella pneumonia*, *Proteus mirabilis*,

Streptococcus faecalis, *Enterobacter species*, *Serratia marcescens*, *Staphylococcus aureus* and *Escherichia coli* by agar well diffusion method. The results demonstrated methanol extract of *Phyllanthus amarus* for highest inhibitory activity against above bacterial species [30]. Similarly, in another study antimicrobial potential of *Phyllanthus amarus* were investigated using agar well diffusion method for activity against several drug resistant pathogens such as *Escherichia coli*, *Pseudomonas aeruginosa*, *Staphylococcus aureus* and *Klebsiella Species*. The results revealed minimum inhibitory concentration (MIC) of the ethanolic plant extracts on *Escherichia coli*, *Staphylococcus aureus*, *Pseudomonas aeruginosa* and *Klebsiella Species* were at 10 mg/ml, 50 mg/ml, 150 mg/ml and 100 mg/ml while the minimum bactericidal concentration were at 50 mg/ml, 100 mg/ml, 150 mg/ml and 150 mg/ml respectively [10]. Further studies on hexane, chloroform, ethyl acetate, acetone and methanol extract of stem bark extracts of *Phyllanthus amarus* demonstrated the antimicrobial activity for all these extracts with a diameter that ranges between 11 mm 24 mm against *Streptococcus pyogenes*, *Staphylococcus aureus*, *Salmonella typhi*, *Escherichia coli*, *Candida albicans*, *Aspergillus flavus* [8]. The antimicrobial activity of the methanolic extract of *Phyllanthus amarus* as studied by agar dilution method and disc diffusion showed significant concentration-dependent antibacterial activity specifically for gram-negative microbes. It was also observed that antibacterial action was mainly due to the isolated phyllanthin [1]. These studies signify the antimicrobial potential of *Phyllanthus amarus* and need of isolation of some potential phytoconstituents from this species.

Antileptospiral Activity:

Leptospirosis is globally important disease found mainly wherever human come in contact with the urine of infected animals or urine contaminated environment. *Phyllanthus amarus* have been investigated for the antileptospiral activity by micro dilution tests and tube dilution technique. The results revealed the inhibitory action of methanolic and aqueous extract of whole plant of *Phyllanthus amarus* against leptospira [31].

Anticonvulsant Activity:

Epilepsy is a major neurological disorder characterized by the occurrence of recurrent seizures. The two widely proposed mechanisms involve

alterations in the voltage-dependent ion channels such as reduction in inhibitory GABA-mediated drive or increase in excitatory glutamate mediated inputs. This chronic progressive CNS disorder affects a large population of the world. In search of herbal treatment, aqueous and ethanolic extract of *Phyllanthus amarus* were evaluated for anticonvulsant effect using pentylenetetrazole (PTZ) and maximal electroshock-induced seizures (MES) in swiss albino rats. The result showed ethanolic and aqueous extract of leaves and stem of *Phyllanthus amarus* significantly effective in abolishing hind limb extension induced by MES as well as PTZ induced seizures [32].

Antidiabetic Activity:

Diabetes is a metabolic disorder of carbohydrate, fat and protein and is considered as the world's largest endocrine disease [33]. The antidiabetic potential of *Phyllanthus amarus* investigated in an experiment model where fasted rats were made diabetic by single intraperitoneal injection of 120 mg/kg of alloxan monohydrates and then two doses of the aqueous and hydroalcoholic extract of *Phyllanthus amarus* administered orally which were then compared with the normal control group that received distilled water only. After 15 days treatment the result demonstrates aqueous and hydroalcoholic extract of *Phyllanthus amarus* decrease the blood glucose level significantly. Serum analysis of the treated experimental animals showed an increase in insulin and reduction in the malondialdehyde concentration, therefore demonstrated the potential antidiabetic property of aqueous and hydroalcoholic extract of *Phyllanthus amarus* [34]. In another study the methanolic extract of *Phyllanthus amarus* was found to inhibit lipid peroxidation & scavenge hydroxyl and superoxide radicals [35]. Since free radicals are linked with diabetes, therefore quenching of free radical could be one mechanism of action [43]. However, there is a need of further experimental studies in order to isolate chemical constituents and their mechanism of action.

Anti-Inflammatory Activity:

The anti-inflammatory potential of *Phyllanthus amarus* was evaluated using different models such as rat Kupffer cells, macrophages RAW264.7, human whole blood and in mice. Two different extracts of *Phyllanthus amarus* (hexane and ethanol/water extracts) and their anti-inflammatory effect was evaluated against the lipopolysaccharide stimulated above mentioned test cells. In addition, anti-inflammatory effect was evaluated in mice that were treated with galactosamine/lipopolysaccharide for inducing acute toxic hepatitis. The evaluation parameters were production of nitrite, prostaglandin E2 and cytokines that were measured by Griess assay, prostaglandin E2 by radioimmunoassay and latter by enzyme-linked immunosorbent assay. The other inflammatory markers such as endotoxin-induced nitric oxide synthase (iNOS) and cyclooxygenase (COX-2) were determined by

Western blot and activation of NF- κ B and activator protein 1 (AP-1) by electrophoretic mobility shift assay (EMSA). The results revealed ethanol/water extracts and hexane extracts effective in inhibition of lipopolysaccharide induced production of nitric oxide (NO) and prostaglandin E2 (PGE2) in Kupffer cells and in macrophages RAW264.7. The extracts also attenuated the lipopolysaccharide induced secretion of tumor necrosis factor (TNF- α) in macrophages RAW264.7 as well as in human whole blood. Hexane and ethanol/water extracts of *Phyllanthus amarus* reduced expression of endotoxin-induced nitric oxide synthase iNOS and cyclooxygenase COX-2 and inhibited activation of nuclear factor NF- κ B. *Phyllanthus amarus* also inhibited induction of interferon- γ (IFN- γ), interleukin (IL)-1 β and interleukin (IL)-10 in human whole blood and reduced tumor necrosis factor (TNF- α) production *in-vivo* [36]. Further, experimental studies have been done to determine the chemical compounds responsible for the activity. In an attempt to test phytoconstituents and extracts of *Phyllanthus amarus* for anti-inflammatory effect, the purified lignans of *Phyllanthus amarus* and different extracts obtained from this plant were evaluated in carrageenan induced paw oedema and neutrophil influx model of inflammation. The result showed that hexane extract and the lignan-rich fraction, or lignansphylltetralin, nirtetralin and niranthin inhibited carrageenan-induced rat paw oedema, lower the increase of interleukin (IL)-1 β tissue levels induced by carrageenan and inhibited neutrophil influx, bradykinin activating factor, platelet activating factor and endothelin-1-induced paw oedema. These results show that the hexane extract, the lignan-rich fraction and the lignansniranthin, phylltetralin and nirtetralin exhibited marked anti-inflammatory properties [37]. Another interesting study where anti-inflammatory effect of soft drink prepared from the leaf extract of *Phyllanthus amarus* was evaluated for its anti-inflammatory effect and the result revealed anti-inflammatory activity of soft drink similar to reference compound Ibuprofen [38]. All these studies acknowledge *Phyllanthus amarus* as potent anti-inflammatory plant and lignins as potent phyto-compounds.

Antifertility Activity:

Phyllanthus amarus possess antifertility activity. This activity was shown in the experimental study where alcoholic extract of *Phyllanthus amarus* brought changes in 3-beta and 17-beta hydroxyl steroid dehydrogenase (HSDs) levels, thereby effecting hormonal conversions in the female mice that confirmed by observation of no pregnancy in cohabited normal females and male mice [39].

Nephroprotective and cardioprotective activity:

Nephroprotective and cardioprotective effect of *Phyllanthus amarus* is evident from the study in which methanol extract of *Phyllanthus amarus* leaves caused a significant dose dependent decrease in the levels of

total cholesterol, urea, total protein, uric acid, and prostatic, alkaline and acid phosphatases, aspartate transaminase (AST) and alanine transaminase (ALT) [40]. Since increase in these enzymes is related to hepatic and heart disorders therefore their reduction shows that the leaves of *Phyllanthus amarus* have hepatoprotective, nephroprotective and cardioprotective properties.

Hepatoprotective effect:

Hepatoprotective effects of aqueous extract from *Phyllanthus amarus* on ethanol-induced rat hepatic injury were studied in *in vitro* study where *Phyllanthus amarus* increases the percentage 3-[4,5-dimethylthiazol-2-yl]-2,5 diphenyltetrazolium bromide (MTT) reduction assay and decreased the release of aspartate transaminase (AST) and alanine transaminase (ALT) in rat primary cultured hepatocytes treated with ethanol. The results reveal that treatment of rats with *Phyllanthus amarus* extract orally brought cell recovery in ethanol-induced liver injury by bringing the levels of aspartate transaminase (AST), alanine transaminase (ALT), high-sensitivity human thyroglobulin (HTG) and Tumor necrosis factor (TNF- α) to normal. Histopathological study confirmed the beneficial effect of *Phyllanthus amarus* with its potential antioxidant activity [41].

Antiviral activity:

Phyllanthus amarus possess antifungal, antiviral and anticancerous properties [4]. Further, evaluation of antiviral activity of *Phyllanthus* species were evident from experiment study where aqueous extract of *Phyllanthus amarus* along with other species of *Phyllanthus* genus were evaluated against Herpes Simplex Virus type-1 and Herpes Simplex Virus type-2 in vero cells by quantitative polymerase chain reaction. Western blot and 2D-gel electrophoresis were used to study protein expressions of treated and untreated infected vero cells. *Phyllanthus amarus* along with *Phyllanthus urinaria* demonstrate the strongest antiviral activity against Herpes Simplex Virus type-1 and Herpes Simplex Virus type-2 which is proposed to its action in the early stage of infection and replication [42].

Haematological Properties:

Phyllanthus amarus has been found to produce some haematological changes in experimental studies. When albino rats were treated with the *Phyllanthus amarus* aqueous extract prepared from the whole plant, dose dependent decrease in erythrocyte sedimentation rate (ESR) and packed cell volume (PCV) was observed. Circulating leucocytes and neutrophils count were significantly increased in rats treated with 100 mg/kg of aqueous extract of *Phyllanthus amarus* as evident by total and differential count studies of blood of experimental animals. In addition quantitative analysis of alanine aminotransferases (ALT) and aspartate aminotransferases (AST) gave significantly higher

values of alanine aminotransferases (ALT) in treated rats. Author has suggested immunostimulant potential of plant [5, 44]

Conclusion:

Phyllanthus amarus herb is widely used Tropical countries including India. It has significant traditional uses, some of them have been experimentally established and an attempt has been made to isolate potential chemical constituents and their mechanism of action. Present review had compiled the traditional uses, pharmacological properties and chemical constituents present, which can be useful information for further study on this plant.

Reference:

- Mazumder A, Mahato A, Mazumder R. Antimicrobial potentiality of *Phyllanthus amarus* against drug resistant pathogens. Natural Product Research 2006; 20(04):323–326.
- Tahseen M, Mishra G. Ethnobotany and Diuretic Activity of Some Selected Indian Medicinal Plants. The Pharma Innovation 2013; 2:112.
- Kumar S, Choudhary H, Seniya C. *In vitro* antibacterial study of aqueous and methanolic extracts of some selected medicinal plants. Journal of Chemical and Pharmaceutical Research 2011; 3:854.
- Ito E, Ukana D, Ekaete D. Phytochemical screening and nutrient analysis of *Phyllanthus amarus*. Asian Journal of Plant Science and Research 2013; 3:116-122.
- Joseph B, Raj SJ. An Overview: Pharmacognostic Property of *Phyllanthus amarus* linn. International Journal of Pharmacology 2011; 1:41.
- Khatoun S, Rai V, Rawat A. Comparative pharmacognostic studies of three *Phyllanthus* species. Journal of Ethnopharmacology 2004; 104:79-86.
- Sen A, Batra A. The study of in vitro and in vivo antioxidant activity and total phenolic content of *Phyllanthus amarus* Schum Thonn: A medicinally important plant. International Journal of Pharmacy and Pharmaceutical Sciences 2013; 5:947.
- Ushie O, Neji P, Etim E. Phytochemical screening and antimicrobial activities of *Phyllanthus amarus* stem bark extracts. International Journal of Modern Biology and Medicines 2013; 3:101-112.
- Chandewar A, Dhongade H. Pharmacognostical and Phytochemical studies of *Phyllanthus amarus* leaves. International Journal of Biomedical and Advance Research 2013; 4:383.
- Adegoke AA, Iberi PA, Akinpelu DA, Aiyegoro P. Studies on phytochemical screening and antimicrobial potentials of *Phyllanthus amarus* against multiple antibiotic resistant bacteria. International Journal of Applied Research in Natural Products 2010; 3:6.
- Patel JR, Tripathi P, Sharma V, Chauhan NS, Dixit VK. *Phyllanthus amarus* Ethnomedicinal uses phytochemistry and pharmacology: A review. Journal of Ethnopharmacology 2011; 138(2):286–313.
- Zingare A. Antimicrobial activity of six members of Euphorbiaceae. New Interdisciplinary National Research Journal 2013; 2:112.
- Naidu MT, Babu NC and Venkaiah M. Medicinal plants used for snakebite in the Kotia hills of Vizianagaram district, Andhra Pradesh. Indian Journal of natural Products and Resources. 2013. 4:194-196.
- Naidu MT, Babu NC, Kumar OA and Venkaiah M. Herbal remedies for rheumatoid arthritis used by the tribes of Vizianagaram district, Andhra Pradesh. Journal of Non-Timber Forest Products. 2012. 19: 303-308.
- Babu NC, Naidu MT and Venkaiah M. Ethnomedicinal plants used by the tribes of Vizianagaram district, Andhra Pradesh. Annals of Pharmaceutical and Pharmacy Science. 2011. 2: 1-4.
- Houghton PJ, Woldemariam TZ, Siobhan OS, Thyagarajan SP. Two securinega type alkaloids from *Phyllanthus amarus*. Phytochemistry 1996; 43:715–717.
- Foo LY. Amariin a di-dehydrohexahydroxydiphenoyl hydrolysable tannin from *Phyllanthus amarus*. Phytochemistry 1993; 33:487–491.
- Foo LY, Wong H, Phyllanthusiin D. an unusual hydrolysable tannin from *Phyllanthus amarus*. Phytochemistry 1992; 31:711–713.
- Kassuya CA, Silvestre A, Menezes-de-Lima Jr O, Marotta DM, Rehder VL, Calixto JB. Antiinflammatory and antiallodynic actions of the lignin niranthin isolated from *Phyllanthus amarus*. Evidence for interaction with platelet activating factor receptor. European Journal of Pharmacology 2006; 546:182-188.
- Srivastava V, Singh M, Malasoni R, Shanker K, Verma RK, Gupta MM *et al.* Separation and quantification of lignans in *Phyllanthus* species by a simple chiral densitometric method. Journal of Separation Science 2008; 31:23-38.
- Leite DF, Kassuya CA, Mazzuco TL, Silvestre A, De-Melo LV, Rehder VL *et al.* The cytotoxic effect and the multidrug resistance reversing action of lignans from *Phyllanthus amarus*. Planta Medica 2006; 72:1353–1358.
- Maciel MAM, Cunha A, Dantas FTNC, Kaiser CR. NMR characterization of bioactive lignans from *Phyllanthus amarus* Schum&Thonn. Journal of Magnetic Resonance Imaging 2007; 6:76–82.
- Londhe JS, Devasagayam TP, Foo LY, Ghaskadbi SS. Radioprotective properties of polyphenols from *Phyllanthus amarus* Linn. Journal of Radiation Research 2009; 50:303–309.
- Foo LY. Amarinic acid and related ellagitannins from *Phyllanthus amarus*. Phytochemistry 1995; 39:217–224.
- Moronkola DO, Ogunwande IA, Oyewole IO, Baser KHC, Ozek T, Ozek G. Studies on the volatile oils of *Momordica charantia* L (Cucurbitaceae) *Phyllanthus amarus* Schum et Thonn (Euphorbiaceae). Journal of Essential Oil Research 2009; 21:393–399.
- Rajeshkumar NV, Joy KL, Kuttan G, Ramsewak RS, Nair MG, Kuttan R. Antitumour and anticarcinogenic activity of *Phyllanthus amarus* extract. Journal of Ethnopharmacology 2002; 81(1):17-22.
- Joshi H, Parle M. Pharmacological evidence for antiamnesic potentials of *Phyllanthus amarus* in mice. African Journal of Biomedical Research 2007; 10:165.
- Lim Y, Murtijaya J. Antioxidant properties of *Phyllanthus amarus* extracts as affected by different drying methods. Food Science and Technology 2007; 40(9):1664-1669.
- Oluwafemi F, Debiri F. Antimicrobial Effect of *Phyllanthus amarus* and *Parquetina nigrescens* on *Salmonella typhi*. African Journal of Biomedical Research 2008; 11(2):215- 219.
- Saranraj P, Sivasakthivelan P. Screening of Antibacterial Activity of the Medicinal Plant *Phyllanthus amarus* Against Urinary Tract Infection Causing Bacterial Pathogens. Applied Journal of Hygiene 2012; 1(3):19-24.
- Chandan S, Umesha S, Balamurugan V. Anti Leptospirosis Antioxidant and DNA damaging properties of *Eclipta alba*

- and *Phyllanthus amarus*. Open Access Scientific Reports. 2012; 1(4):1-8.
32. Manikkoth S, Deepa B, Joy AE, Rao S. Anticonvulsant activity of *Phyllanthus amarus* in experimental animal models 2011; 4:144-149.
33. Moshi MJ, Lutale JJ, Rimoy GH, Abbas ZG, Josiah RM, Swai AB. The effect of *Phyllanthus amarus* aqueous extract on blood glucose in non-insulin dependent diabetic patients. Phytotherapy Research 2001; 15(7):577-80.
34. Evi PL, Degbeku K. Antidiabetic Activity of *Phyllanthus amarus* Schum and Thonn on Alloxan induced diabetes in Male Wistar Rats. Journal of Applied Sciences 2011; 11(16):2968-2973.
35. Kiran D, Rohilla A, Rohilla S. *Phyllanthus amarus* an ample therapeutic potential herb. International Journal of Research in Ayurveda & Pharmacy 2011; 2(4):1099.
36. Kierner AK, Hartung T, Huber C, Vollmar AM. *Phyllanthus amarus* has anti-inflammatory potential by inhibition of iNOS, COX-2 and cytokines via the NF- κ B pathway. Journal of Hepatology 2003; 38(3):289-97.
37. Kassuya CA, Leite DF, De-Melo LV, Rehder VL, Calixto JB. Anti-inflammatory properties of extracts, fractions and lignans isolated from *Phyllanthus amarus*. Planta Medica 2005; 71(8):721-6.
38. Adeolu AA, Sunday OO. Antiinflammatory and analgesic activities of soft drink leaf extract of *Phyllanthus amarus* in some laboratory animals. British Biotechnology Journal 2013; 3:191-204.
39. Rao MV, Alice KM. Contraceptive effects of *Phyllanthus amarus* in female mice. Phytotherapy Research 2001; 15(3):265-7.
40. Obianime AW, Uchie FI. The phytochemical screening and the effects of methanolic extract of *Phyllanthus amarus* leaf on the biochemical parameters of male guinea pigs. Journal of Applied Sciences and Environmental Management 2008; 12(4):73-77.
41. Pramyothin P, Ngamtin C, Pongshompoo S, Chaichantipyuth C. Hepatoprotective activity of *Phyllanthus amarus* Schum Thonn extract in ethanol treated rats: In vitro and in vivo studies. Journal of Ethnopharmacology 2007; 114(2):169-173.
42. Tan W, Jaganath I, Manikam I. Evaluation of antiviral activities of four local Malaysian *Phyllanthus* species against Herpes simplex viruses and possible antiviral target. International Journal of Medical Sciences 2013; 10(13):1817-1892.
43. Oberley LW. Free radicals and diabetes. Free Radical Biology and Medicine 1988; 5(2):113-24.
44. Taiwo IA, Oboh BO, Francis-Garuba PN. Haematological properties of aqueous extract of *Phyllanthus amarus* (Schum and Thonn.) and *Xylopi aethiopica* (Dunal) a rich in albino rats. Ethno-Med 2009; 3(2):99-103.